



Bright Futures Parent Handout 9 and 10 Year Visits

Here are some suggestions from Bright Futures experts that may be of value to your family.

NUTRITION AND PHYSICAL ACTIVITY

Staying Healthy

- Encourage your child to eat healthy.
- Buy fat-free milk and low-fat dairy foods, and encourage 3 servings each day.
- Include 5 servings of vegetables and fruits at meals and for snacks daily.
- Limit TV and computer time to 2 hours a day.
- Encourage your child to be active for at least 1 hour daily.
- Eat as a family often.

SAFETY

Safety

- The back seat is the safest place to ride in a car until your child is 13 years old.
- Use a booster seat until the vehicle's safety belt fits. The lap belt can be worn low and flat on the upper thighs. The shoulder belt can be worn across the shoulder and the child can bend at the knees while sitting against the vehicle seat back.
- Teach your child to swim and watch her in the water.
- Your child needs sunscreen (SPF 15 or higher) when outside.
- Your child needs a helmet and safety gear for biking, skating, in-line skating, skiing, snowmobiling, and horseback riding.
- Talk to your child about not smoking cigarettes, using drugs, or drinking alcohol.
- Make a plan for situations in which your child does not feel safe.
- Get to know your child's friends and their families.
- Never have a gun in the home. If necessary, store it unloaded and locked with the ammunition locked separately from the gun.

DEVELOPMENT AND MENTAL HEALTH

Your Growing Child

- Be a model for your child by saying you are sorry when you make a mistake.
- Show your child how to use his words when he is angry.
- Teach your child to help others.
- Give your child chores to do and expect them to be done.
- Give your child his own space.
- Still watch your child and your child's friends when they are playing.
- Understand that your child's friends are very important.
- Answer questions about puberty.
- Teach your child the importance of delaying sexual behavior. Encourage your child to ask questions.
- Teach your child how to be safe with other adults.
 - No one should ask for a secret to be kept from parents.
 - No one should ask to see your child's private parts.
 - No adult should ask for help with his private parts.

SCHOOL

School

- Show interest in school activities.
- If you have any concerns, ask your child's teacher for help.
- Praise your child for doing things well at school.
- Set a routine and make a quiet place for doing homework.
- Talk with your child and her teacher about bullying.

ORAL HEALTH

Healthy Teeth

- Help your child brush teeth twice a day.
 - After breakfast
 - Before bed
- Use a pea-sized amount of toothpaste with fluoride.
- Help your child floss his teeth once a day.
- Your child should visit the dentist at least twice a year.
- Encourage your child to always wear a mouth guard to protect teeth while playing sports.

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™


The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.

10 tips

Nutrition
Education Series

build a healthy meal

10 tips for healthy meals



A healthy meal starts with more vegetables and fruits and smaller portions of protein and grains. Think about how you can adjust the portions on your plate to get more of what you need without too many calories. And don't forget dairy—make it the beverage with your meal or add fat-free or low-fat dairy products to your plate.

1 make half your plate veggies and fruits

Vegetables and fruits are full of nutrients and may help to promote good health. Choose red, orange, and dark-green vegetables such as tomatoes, sweet potatoes, and broccoli.

2 add lean protein

Choose protein foods, such as lean beef and pork, or chicken, turkey, beans, or tofu. Twice a week, make seafood the protein on your plate.



3 include whole grains

Aim to make at least half your grains whole grains. Look for the words "100% whole grain" or "100% whole wheat" on the food label. Whole grains provide more nutrients, like fiber, than refined grains.

4 don't forget the dairy

Pair your meal with a cup of fat-free or low-fat milk. They provide the same amount of calcium and other essential nutrients as whole milk, but less fat and calories. Don't drink milk? Try soy milk (soy beverage) as your beverage or include fat-free or low-fat yogurt in your meal.



5 avoid extra fat

Using heavy gravies or sauces will add fat and calories to otherwise healthy choices. For example, steamed broccoli is great, but avoid topping it with cheese sauce. Try other options, like a sprinkling of low-fat parmesan cheese or a squeeze of lemon.

6 take your time

Savor your food. Eat slowly, enjoy the taste and textures, and pay attention to how you feel. Be mindful. Eating very quickly may cause you to eat too much.

7 use a smaller plate

Use a smaller plate at meals to help with portion control. That way you can finish your entire plate and feel satisfied without overeating.

8 take control of your food

Eat at home more often so you know exactly what you are eating. If you eat out, check and compare the nutrition information. Choose healthier options such as baked instead of fried.

9 try new foods

Keep it interesting by picking out new foods you've never tried before, like mango, lentils, or kale. You may find a new favorite! Trade fun and tasty recipes with friends or find them online.



10 satisfy your sweet tooth in a healthy way

Indulge in a naturally sweet dessert dish—fruit! Serve a fresh fruit cocktail or a fruit parfait made with yogurt. For a hot dessert, bake apples and top with cinnamon.

VACCINE INFORMATION STATEMENT

Tdap Vaccine

What You Need to Know

(Tetanus,
Diphtheria and
Pertussis)

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Tetanus, diphtheria and pertussis are very serious diseases. Tdap vaccine can protect us from these diseases. And, Tdap vaccine given to pregnant women can protect newborn babies against pertussis.

TETANUS (Lockjaw) is rare in the United States today. It causes painful muscle tightening and stiffness, usually all over the body.

- It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of 10 people who are infected even after receiving the best medical care.

DIPHThERIA is also rare in the United States today. It can cause a thick coating to form in the back of the throat.

- It can lead to breathing problems, heart failure, paralysis, and death.

PERTUSSIS (Whooping Cough) causes severe coughing spells, which can cause difficulty breathing, vomiting and disturbed sleep.

- It can also lead to weight loss, incontinence, and rib fractures. Up to 2 in 100 adolescents and 5 in 100 adults with pertussis are hospitalized or have complications, which could include pneumonia or death.

These diseases are caused by bacteria. Diphtheria and pertussis are spread from person to person through secretions from coughing or sneezing. Tetanus enters the body through cuts, scratches, or wounds.

Before vaccines, as many as 200,000 cases of diphtheria, 200,000 cases of pertussis, and hundreds of cases of tetanus, were reported in the United States each year. Since vaccination began, reports of cases for tetanus and diphtheria have dropped by about 99% and for pertussis by about 80%.

2 Tdap vaccine

Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. One dose of Tdap is routinely given at age 11 or 12. People who did *not* get Tdap at that age should get it as soon as possible.

Tdap is especially important for healthcare professionals and anyone having close contact with a baby younger than 12 months.

Pregnant women should get a dose of Tdap during **every pregnancy**, to protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Another vaccine, called Td, protects against tetanus and diphtheria, but not pertussis. A Td booster should be given every 10 years. Tdap may be given as one of these boosters if you have never gotten Tdap before. Tdap may also be given after a severe cut or burn to prevent tetanus infection.

Your doctor or the person giving you the vaccine can give you more information.

Tdap may safely be given at the same time as other vaccines.

3 Some people should not get this vaccine

- A person who has ever had a life-threatening allergic reaction after a previous dose of any diphtheria, tetanus or pertussis containing vaccine, OR has a severe allergy to any part of this vaccine, should not get Tdap vaccine. Tell the person giving the vaccine about any severe allergies.
- Anyone who had coma or long repeated seizures within 7 days after a childhood dose of DTP or DTaP, or a previous dose of Tdap, should not get Tdap, unless a cause other than the vaccine was found. They can still get Td.
- Talk to your doctor if you:
 - have seizures or another nervous system problem,
 - had severe pain or swelling after any vaccine containing diphtheria, tetanus or pertussis,
 - ever had a condition called Guillain-Barré Syndrome (GBS),
 - aren't feeling well on the day the shot is scheduled.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4 Risks

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own. Serious reactions are also possible but are rare.

Most people who get Tdap vaccine do not have any problems with it.

Mild problems following Tdap (*Did not interfere with activities*)

- Pain where the shot was given (about 3 in 4 adolescents or 2 in 3 adults)
- Redness or swelling where the shot was given (about 1 person in 5)
- Mild fever of at least 100.4°F (up to about 1 in 25 adolescents or 1 in 100 adults)
- Headache (about 3 or 4 people in 10)
- Tiredness (about 1 person in 3 or 4)
- Nausea, vomiting, diarrhea, stomach ache (up to 1 in 4 adolescents or 1 in 10 adults)
- Chills, sore joints (about 1 person in 10)
- Body aches (about 1 person in 3 or 4)
- Rash, swollen glands (uncommon)

Moderate problems following Tdap (*Interfered with activities, but did not require medical attention*)

- Pain where the shot was given (up to 1 in 5 or 6)
- Redness or swelling where the shot was given (up to about 1 in 16 adolescents or 1 in 12 adults)
- Fever over 102°F (about 1 in 100 adolescents or 1 in 250 adults)
- Headache (about 1 in 7 adolescents or 1 in 10 adults)
- Nausea, vomiting, diarrhea, stomach ache (up to 1 or 3 people in 100)
- Swelling of the entire arm where the shot was given (up to about 1 in 500).

Severe problems following Tdap (*Unable to perform usual activities; required medical attention*)

- Swelling, severe pain, bleeding and redness in the arm where the shot was given (rare).

Problems that could happen after any vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at fewer than 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious problem?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.
- Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your doctor. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement Tdap Vaccine

2/24/2015

42 U.S.C. § 300aa-26

